



Mapping Terrestrial Groundwater Dependent Ecosystems:

Method Development and Example Output

P. Evan Dreisel
Rob Clark
Xiang Cheng
Mark Reid
Alister Terry
Jonathon Fawcett
Denise Cochrane

If you would like to receive this information/publication in an accessible format (such as large print or audio) please call the Customer Service Centre on 136 186, TTY 1800 122 969, or email customer.service@dpi.vic.gov.au.

Citation:

Dresel, P. E., Clark, R. Cheng, X., Reid, M., Fawcett, J., and Cochraine, D. (2010) Mapping Terrestrial Groundwater Dependent Ecosystems: Method Development and Example Output. Victoria Department of Primary Industries, Melbourne VIC. 66 pp.

ISBN 978-1-74264-321-2 (print)
ISBN 978-1-74264-322-9 (CD-ROM)
ISBN 978-1-74264-323-6 (online)

Published by the Department of Primary Industries
, April 2010

© The State of Victoria 2010.

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the Copyright Act 1968.

Authorised by the Department of Primary Industries
1 Spring Street, Melbourne 3000.

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

For more information about DPI go to www.dpi.vic.gov.au or phone the Customer Service Centre on 136 186.

Contents

Figures	4
Tables	5
Acronyms and Definitions	6
1 Introduction	9
1.1 Project Objectives	9
2 What are Groundwater Dependent Ecosystems?	11
2.1 Original Classification System from Hatton & Evans (1998)	11
2.2 Additional Classes of GDEs	11
2.3 Alternative GDE Classification System by Eamus et al. (2006)	12
2.4 ADE classification & identification by Colvin et al. (2007)	12
3 Description of GDE Classes	14
3.1 Terrestrial vegetation	14
3.2 Wetlands - fresh and saline groundwater dependency	15
3.3 Coastal estuarine and near shore marine systems	17
3.4 River base flow systems	18
3.5 Aquifer and cave ecosystems	18
3.6 Terrestrial fauna	18
4 Known Victorian areas with GDEs	20
5 Identification of GDEs	21
5.1 Vegetation analysis	21
5.2 Groundwater – surface water interaction	22
5.3 Remote sensing approaches	23
5.4 Geological mapping approaches	23
6 Development of state-wide potential GDE maps	25
6.1 Spatial data sources for GDE mapping	25
6.2 Definition of Aridity Zones	27
6.3 Remote sensing methodology	28
6.3.1 Contrasting mid-summer photosynthetic activity	28

6.3.2	Consistent annual photosynthetic activity	29
6.3.3	Unsupervised classification of Landsat spectral data	33
6.4	Integration of data sets and development of Model layers	33
6.5	Classification and Attribution of GDEs	37
6.5.1	Groundwater Interactive Map	38
6.5.2	Geomorphologic, Geologic, and Chemical Attributes	39
7	CMA region potential GDE map results	42
7.1	Mallee CMA	42
7.2	Wimmera CMA	42
7.3	Glenelg Hopkins CMA	42
7.4	North Central CMA	46
7.5	Corangamite CMA	48
7.6	Port Phillip and Westernport CMA	50
7.7	Goulburn Broken CMA	50
7.8	West Gippsland CMA	50
7.9	North East CMA	50
7.10	East Gippsland CMA	55
8	Conclusions and Recommendations	58
8.1	Limitations of the maps	58
8.2	Recommended usage of the maps in current form	59
9	References	61

Figures

Figure 1.	Distribution of wetlands where previous research has established a link with groundwater and location of stream gauges where base flow calculations have been made	20
Figure 2.	Catchment Management Authority (CMA) regions for Victoria	25
Figure 3.	Climate zones for the Corangamite CMA region derived from aridity indices (A) and comparing the climate zones to the MODIS EVI time series data for summer 2003 (B)	27
Figure 4.	Process flow for development of potential terrestrial GDE map layer	28
Figure 5.	Mid summer NDVI for the combined years of 1995 and 2002 in the North Central CMA region	30
Figure 6.	Time series of MODIS EVI values showing consistent annual photosynthetic activity of redgum forest in comparison to variable annual photosynthetic activity of pasture	31
Figure 7.	EVI standard deviation layer for the North Central CMA region, showing the 27 classes with the lowest standard deviation	32
Figure 8.	Distribution of selected EVC groupings used in the unsupervised classification in Victoria. Black outline defines the Glenelg-Hopkins CMA region.	34
Figure 9.	Application of land use mask to potential terrestrial GDEs. A. Land use classification for section of North East CMA region. B. Potential terrestrial GDEs before and after application of mask.	36
Figure 10.	Indicative groundwater interactive map (GIM) for the North Central CMA region showing the location of piezometers that have recorded a watertable depth of less than 5 m. Colour coding shows five different percentile	

	ranges indicating percentage of piezometer record for which the water table was less than 5 m below ground surface	40
Figure 11.	Potential terrestrial GDEs in the Mallee CMA. A. Overview. B. GDEs mapped in area of extensive mallee vegetation. C. Sparse association of mapped GDEs with lakes, salinas and areas of shallow water table	43
Figure 12.	Potential terrestrial GDEs in the Wimmera CMA. A. Overview. B. Area of known groundwater-surface water interaction near Edenhope. C. Area of potential GDEs near the Wimmera River at Dimboola	44
Figure 13.	Potential terrestrial GDEs in the Glenelg Hopkins CMA. A. Overview. B. Detail of GDEs shown around the southern Grampians. C. Potential GDEs and spring locations northwest of Portland.	45
Figure 14.	Map of the Cockajemmi Lakes region showing potential GDEs and the extent of the Quaternary age basalt flow	46
Figure 15.	Potential GDE map for the North Central CMA region. A. Overview of the CMA region. B. Area between Swan Hill and Kerang showing the association of potential GDEs in areas where surface water drainages or wetlands intersect groundwater interactive landscapes. C. Relation of GDEs to spring locations in the vicinity of Daylesford.	47
Figure 16.	Map of potential GDEs for the Tarpaulin Creek area, showing landscapes that may contain GDEs with shallow rooted vegetation (cyan) and landscapes which might include deep rooted vegetation and fall outside the indicative groundwater interaction map (yellow)	48
Figure 17.	Potential terrestrial GDEs in the Corangamite CMA. A. Overview. B. Detail showing areas of potential groundwater dependency east and south of Ballarat. C. Potential terrestrial GDEs around lakes near Colac, including Lake Corangamite	49
Figure 18.	Potential terrestrial GDEs in the Port Phillip and Westernport CMA. A. Overview. B. Detail showing areas of potential groundwater dependency along streams near Bacchus Marsh. C. Potential GDEs in the Yarra River riparian zone near Healesville	51
Figure 19.	Satellite image showing potential terrestrial GDEs along Yarra River	52
Figure 20.	Potential terrestrial GDEs in the Goulburn Broken CMA. A. Overview. B. Detail of GDEs in the Murray Riverfloodplain (large green area is the Barmah Forest), along the lower Goulburn River (lower left) and along other streams including the Broken Creek	53
Figure 21.	Potential terrestrial GDEs in the West Gippsland CMA. A. Overview. B&C. Detail showing areas of potential groundwater dependency along streams and waterways in the Latrobe Valley (B) and north of Lake Wellington (C)	54
Figure 22.	Potential terrestrial GDEs in the North East CMA. A. Overview. B. Detail showing areas of potential groundwater dependency along the Murray and Ovens River floodplains	55
Figure 23.	Potential terrestrial GDEs in the East Gippsland CMA. A. Overview. B. Detail showing potential groundwater dependency in areas surrounding drainage lines and water bodies and other scattered areas. The large blocks are responding to open water	57
Figure 24.	Potential terrestrial GDEs in a section of the North East CMA region, classified by groundwater interactive landscape. The areas of greatest interest are expected to be in the shallow water table zones	60

Tables

Table 1	Some references of some research into groundwater-surface water interaction and groundwater use by vegetation within Victorian and adjacent landscapes	20
Table 2.	Tools to identify GDE's (after Clifton et al. 2007)	22
Table 3.	EVC groups and examples used to develop the indicative response layer	33
Table 4.	Example weighting factors for use in model combining remote sensing GDE classes	35
Table 5.	Land use classes used to exclude areas from classification as potential terrestrial GDEs	37
Table 6.	Attributes assigned to each potential GDE location	41

Acronyms and Definitions

AGO	Australian Greenhouse Office
CGDL	State of Victoria Corporate Geospatial Data Library
CMA	Catchment Management Authority
EVC	Ecological Vegetation Class
EVI	Enhanced Vegetation Index
GDE	Groundwater Dependent Ecosystem
GFS	Groundwater Flow System
GIM	Groundwater Interactive Map
GMU	Geomorphologic Management Unit
NDVI	Normalised Difference Vegetation Index

