

Victorian Weed Risk Assessment (WRA) method

Methodology	2
1.1 Decision Support Systems	2
1.2 Applying the Analytic Hierarchy Process to weed risk assessment	3
1.3 Weed Risk Assessment in Victoria	4
1.4 Invasiveness Potential of Pest Plants	7
1.5 Present and Potential Distribution of Pest Plants	15
1.6 Present Distribution	15
1.7 Potential Distribution	18
1.8 Ratio of Present to Potential Distribution	20
1.9 Limitations of Present and Potential Distribution Maps	22
1.10 Impacts	23
1.11 Calculating a weed risk score	24
References	32
Appendix 1. Invasiveness criteria and intensity ratings	34
Appendix 2. Impact criteria and intensity ratings	38

Methodology

1.1 Decision Support Systems

Natural resource managers work with complex systems where problem solving and decision-making is based on extensive, but incomplete, uncertain, and even contradictory data and knowledge. There is often no single correct method, or answer, to address problems in these systems. Managers therefore require a decision making process to break down complex systems into simpler steps with defined criteria to allow assessment and prioritisation of issues.

We propose a specific decision support system (DSS) that relies on expert quantitative and qualitative data. This DSS relies on a type of multi-criteria analysis (analytic hierarchy process or AHP) that enables complex issues to be broken down into sets of related criteria. The AHP (Saaty 1995) is a method that assists with decisions about priorities using qualitative and/or quantitative information. It facilitates effective decisions on complex issues by simplifying and expediting the intuitive decision-making process. AHP does this by organising a complex unstructured situation into component parts with similar themes, arranges these parts into a hierarchical order, assigns values relative to each variable, and synthesises these judgements to determine which variables are most important. AHP also provides an effective structure for group decision-making. This is generally based either on already documented scientific information or in workshop sessions with experts.

Because there is often a lack of specific information on land and resource value, and the impact of any particular weed on social, environmental and economic resources, there is a need for a methodology that considers qualitative and quantitative information. The DSS allows for this integration and applies visible weighting to criteria or resources to indicate their importance. A summary of the analytic hierarchy process, as described by Saaty (1995), is presented in Table 1.

The main benefits of using this type of decision support system are that:

- It takes advantage of existing information by integrating it into a system that allows a wide range of users to interpret the data, using a methodology developed by experts.
- It captures the expertise of specialists and makes this expertise available across a wide range of decision-making contexts.
- It provides an explicit method for integrating ecological, social, and economic criteria into the decision-making process.
- It can provide a set of "best practice" decision-making tools to planners and managers.
- It provides a mechanism for identifying information shortfalls.
- It enables a qualitative analysis of the suitability of data and its relevance to the decision-making process.
- It provides a framework for developing sophisticated benchmarks, including identifying the necessary trade-offs between competing value systems.
- It is easily up-dated as research fills knowledge gaps.

Table 1. Analytic hierarchy process steps as described by Saaty (1995).

SAATY	
HIERARCHY	
1	Define the problem and specify the solution desired
2	Structure the hierarchy
WEIGHTING	
3	Construct a pair-wise comparison matrix
4	Obtain all judgements required to develop the set of matrices
5	Test consistency
6	Perform 3 – 5 for all levels and clusters in the hierarchy
7	Use hierarchical composition to weight the vectors of priorities by the weights of the criteria, and take the sum over all weighted priority entries corresponding to those in the next lower level and so on.
Evaluate the consistency of the entire hierarchy.	

Source: "Priority setting Framework for Natural Resources Management – Application of the Analytical Hierarchical Process and Natural Resources Accounting" (Sposito *et al* 2002)

To be scientifically valid though, any system developed must meet certain criteria:

- 1) "It must be transparent, be open to review, and have been evaluated by peers.
- 2) It must have a logical framework that includes independent factors-identified through critical observation, experimentation, or both-important in the invasion process.
- 3) Use of the framework must be repeatable and lead to the same outcome, regardless of who makes the predictions." (NRC 2002)

Although the US National Research Council (NRC 2002) applied these criteria specifically to systems predicting invasiveness, they should apply equally to all components of a decision support system.

1.2 Applying the Analytic Hierarchy Process to weed risk assessment

The species that are of highest risk are those that have the greatest potential to affect valued resources. However the degree of affect can only be determined if managers responsible for those resources prioritise or value them in relation to each other. This process can be accomplished through workshops using the AHP – DSS to rank the social, environmental and economic resources of the region. Any process developed for a territory or State in Australia though should address the requirements of the Australian Standard for National Post-Border Weed Risk Management (AS/NZS HB 294:2006 Standards Australia/Standards New Zealand 2006); the method described here meets these requirements.

The information that is needed to enable threats to be assessed under this process includes:

- The species that could threaten the region either now or in the future.
- Information about the biology of each species and its potential rate of spread.
- The level of impact that a species could have on social, agricultural and environmental resources.
- The values that land managers assign to affected resources.

With this information, the relative importance of invasive species can be determined by considering:

- 1) How invasive it is, i.e., how fast can the species spread. Generally this relates to the intrinsic biological features of the species (i.e. dispersal, reproductive and competitive rate).
- 2) The present and potential extent of the species.
- 3) And importantly, what social, environmental, and economic impacts the species has and the value of the things that are impacted upon.

1.3 Weed Risk Assessment in Victoria

To make informed decisions about the best way to control weeds on public land in Victoria, it is necessary that the relative importance of each weed be determined. It is essential that this is done prior to the allocation of priority works or funding. The Australian Standard for National Post-Border Weed Risk Management Protocol (AS/NZS HB 294:2006 Standards Australia/Standards New Zealand, 2006), for example, states that "a semi-quantitative analysis is the most appropriate for ranking species where there are considerable, long-term financial investments in weed management". Decisions based on limited factual data and emotional reactions will almost certainly result in unnecessary expenditure of resources and damage to the environment through inappropriate use of control measures. Consider the situation in Victoria, where over 1200 plant species are naturalised or incipiently naturalised (Ross and Walsh 2003). It has been estimated that only about ten percent of naturalised plant species become weeds of significant economic and ecological impact (Williamson and Fitter 1996). It is therefore unrealistic and unnecessary to expect that all alien plants can, and should, be controlled.

The Weed Risk Assessment (WRA) developed by the Biosciences Research Division of the Department of Primary Industries, Victoria, is a prioritisation process or risk assessment, based on the AHP, which ranks weeds by:

- 1) Assessing the plant's invasiveness.
- 2) Comparing the plant's present and potential distribution; and
- 3) Determining the impacts of the plant on social, economic, and environmental values.

The WRA is therefore expressed as a hierarchy (Figure 1), the components of which are weighted (using AHP) to allow the determination of a weed risk score for individual species.

The weed risk score is expressed as:

$$\text{Weed risk score} = \alpha (\text{Invasiveness score}) + \beta (\text{Present:Potential Distribution}) + \delta (\text{Impact})$$

(where α , β and δ are weightings).




Figure 1. Hierarchy illustrating components of the Weed Risk Assessment.

1.3.1 Assumptions

No specific targeted control

For each criterion (both invasiveness and impact), species are assessed on their potential in the absence of targeted control (e.g., no change in routine herbicide use to specifically target the weed of interest). Targeted control is a consequence of a weed being assessed as a significant threat.

Limited information on species

To assess plants for both invasiveness and impact, information from a variety of sources including databases, journal articles, floras of the world (books or articles describing the species of a particular country or region), online information, and other sources was accessed. However, information relating directly to specific criteria is not always available. Where such information is lacking, there are two options; rate the criterion as Medium (M) or, where suitable other information is available, estimate a likely response. By assigning a rating of M the maximum possible error is ± 0.5 for that criterion. Assigning a rating of H or L could introduce an error of ± 1 .

In some cases an answer can be implied from other information about the plant. For example, a weedy grass would be considered to contribute to an increase in fire frequency (though not intensity) due to, say, its documented ability to dominate its environment and suppress (less fire-prone) herbaceous species. There may be no specific mention of the plant's ability to change the fire regime, but in this case we could confidently score the criterion as Medium Low (ML) rather than applying the Medium score.

Degree of affect

Plants are assessed for their potential to affect natural or agricultural landscapes negatively. The rating chosen is based on the assumption that a plant will achieve its maximum growth and/or impact. For example, Paterson's curse (*Echium plantagineum* L.) is often regarded as non-toxic, yet research has shown that toxic principles within the plant can cause liver damage sometimes leading to animal death (Parsons & Cuthbertson 2001). So, while experience suggests the plant is harmless, there is evidence that indicates otherwise. Accordingly, this species is rated as being toxic to native fauna. It is given a MH rating, rather than H, based on the presumption that native fauna will be able to browse on a variety of species, not solely Paterson's curse.

We acknowledge that a species will not always find optimal conditions in every situation, but it is the only way of consistently assessing a range of plants.

This risk assessment process is generic. It enables a large number of species to be evaluated in a short time and to be ranked according to the score each plant achieves. The assessment of any one plant only has relevance to the other plants assessed, it does not confer any inherent qualities, either good or bad, about the plant. The results are used to compare assessed species and rank, or prioritise them accordingly.

1.3.2 Rationale in weighting Invasiveness, Distribution and Impact

Researchers of the Cooperative Research Centre for Australian Weed Management (Weeds CRC) and Department of Primary Industries (DPI) weed experts determined a preliminary ranking of the three subcomponents of the WRA. The basis of the weighting was that invasiveness was considered less important than distribution, which in turn was considered less important than impact, with the following ratios:

- Invasiveness is 3 time less important than distribution
- Invasiveness is 4 times less important than impact
- Distribution is half as important than impact

A preliminary AHP pair-wise comparison produced the following weightings (with an acceptable consistency ratio of 0.02) for invasiveness, distribution and impact:

Invasiveness - 12% Distribution - 32% Impact - 56%

Therefore, when calculating a weed risk score; $\alpha = 0.12$, $\beta = 0.32$ and $\delta = 0.56$

The method for developing scores for each of the subcomponents; invasiveness, present and potential distribution, and impact, is outlined in Sections 1.4-1.6).

1.3.3 Confidence

As noted in Assumptions (Section 1.3.1 above) an absence of information can be treated in two ways, either infer from other data or score the criterion as medium. In either case, the lack of absolute information casts immediate doubt on the accuracy of the response. A refinement to that approach, which can be applied to all criteria and thus to the complete assessment, is that of a confidence rating for each answer. The confidence rating is based on the quality of reference material(s) used to answer a question. This approach follows the method used by Robertson *et al* (2003), which indicates uncertainty and availability of data for each criterion. The lower the confidence score the greater the uncertainty and amount of missing data for that criterion. This approach has the advantage that it explicitly indicates a level of confidence in the total risk score assigned to a species. That is, it can be used as a measure of how much faith should be placed in a given risk score, and that further research is desirable. In addition, the confidence score can be used as a measure of the state of knowledge of a given species. Intensity ratings (ie. typical information sources and their relative quality rating) for the confidence scores are listed in Table 2 below.

Table 2. Confidence score intensity ratings

Document Type Or Information Source	Rating	Score
<ul style="list-style-type: none"> Peer-reviewed scientific paper 	H	1
<ul style="list-style-type: none"> High quality science or plant specific books (eg. floras), Non-peer reviewed scientific paper (eg. conference proceedings), Personal communications from expert (eg. PhD, or higher degree on species being assessed), Unpublished reports from highly reliable source (eg. commercial reports or honours theses, etc.), Internet information from Herbaria data, or Internet information that cites sources from MH category, as listed above. 	MH	0.75
<ul style="list-style-type: none"> Personal communications from people with experience with the species under assessment, Information from general plant books (eg. Encyclopaedia Botanica, Gardening Flora, etc.), Unpublished reports from uncertain sources, Internet information that cites sources from M category, or Internet information from government or university websites (eg. Australian state governments, or USDA) 	M	0.5
<ul style="list-style-type: none"> Anecdotal data from non-experts, Internet information that cites anecdotal non-expert sources, Internet information from uncertain/uncited sources, or Horticultural, nursery notes or general web pages. 	ML	0.25
<ul style="list-style-type: none"> No data or reference material available. 	L	0

The assessment confidence score is calculated by giving equal weighting to the confidence score for each question, and then adding them together to give a total between 0 and 1. Where information relating directly to specific criteria is not available, the risk rating assigned is generally medium (M) with a correspondingly low confidence level.

By comparing the confidence score for each species with the Confidence score intensity ratings, you will gain an understanding of the standard of literature available in general for that species. For example, *Acacia longifolia* has a confidence score of 0.62. This indicates that on average, the quality of the literature for this species was between M and MH. Although some questions would have used high quality (H) data, and others no information (L), the standard of literature was generally better than information from general plant books and unpublished reports, but not as good a quality as conference proceedings or personal communications with species experts.

1.4 Invasiveness Potential of Pest Plants

Many researchers have focused on the relative invasiveness of species as an indicator of potential spread rate. Invasiveness can be defined as the ability to establish, reproduce, and disperse within an

ecosystem. Plant propagules arrive at a new site with certain inherent characteristics that previously enabled their successful survival and continued reproduction throughout their evolutionary history. There is no single suite of characteristics which make a plant invasive, rather there are several predisposing factors that act either alone or together to increase the chance of a plant becoming invasive.

Many researchers have also agreed that the following biological attributes of a plant species are associated with invasiveness.

- *Ecological status; a generalist or specialist plant.*
Most common and noxious weeds in southern Australia are generalist and opportunistic rather than requiring specific niches or special habitat requirements.
- *'Weedy' phenology and biology; such as competitive growth, seed dispersal mechanisms, seed dormancy and propagule production.*
Major weeds can have attributes such as high seed production, rapid vegetative spread, long-lived seeds, staggered germination, competitive growth and long-distance seed dispersal. However, there is no defined group of ecological and biological attributes that can be used to identify all major weeds. Different attributes may be important for different plant families and different ecosystems.
- *Wide native range.*
Within a genus the more important weeds may have a wider native range.
- *Taxonomic position; members of generally 'weedy' plant families.*
Certain plant families such as Poaceae (grasses), Asteraceae (eg. daisies, thistles), Iridaceae (irises) and Brassicaceae (eg. mustards, turnips) are noted for having many 'weedy' species.
- *Effective modes of reproduction and genetic variation.*
Plant species that vegetatively reproduce or self-pollinate have the potential to start new populations from a single, isolated plant. However, high levels of inbreeding in self-pollinators may limit their adaptability compared to cross-pollinators.

Other factors may also favour invading species. For example invading species are generally free of the biotic interactions that occur in their natural range, providing them with a competitive advantage over native species that have many co-evolved predators present (Sax & Brown 2000). As this is not a specific biological attribute of a plant it has not been included in the invasiveness assessment criteria.

Specific criteria for a generic model to assess the potential invasiveness of weeds were determined at two national workshops, held at the Arthur Rylah Institute (ARI), in June 1998. A working party at the Department of Primary Industries in Frankston (previously the Keith Turnbull Research Institute) then used an expert system, relying on multi-criteria analysis/analytical hierarchical process (AHP) (Saaty 1995), to develop a decision-tree that allows groups and criteria to be weighted according to importance (

Table 3).

These criteria are echoed in the Australian Standard for National Post-Border Weed Risk Management Protocol (AS/NZS HB 294:2006 Standards Australia/Standards New Zealand, 2006). The protocol was developed to foster the use and further development of decision support systems for prioritising weed species for management at the regional, state/territory and national levels. That document nominates the following attributes as important measures by which to assess invasiveness.

- What is the weed's ability to establish amongst existing vegetation?
- What is the weed's reproductive ability? This includes attributes of time to seeding, seed production and vegetative reproduction
- What is the weed's dispersal ability? This incorporates wind, water, flying animals, ground animals, deliberate human spread, accidental human spread, vehicles and produce or byproduct contaminants.

The criteria used in this process expands upon the attributes nominated in the proposed national specification.

Table 3. Group and criteria weightings for determining invasive potential.

GROUP	CRITERIA	GROUP & CRITERIA WEIGHTINGS	TOTAL CRITERIA WEIGHTINGS
Establishment		0.500	
	Germination requirements?	0.085	0.0425
	Establishment requirements?	0.671	0.3355
	Disturbance requirements?	0.244	0.122
Growth/competitive ability		0.096	
	Life form?	0.060	0.00576
	Allelopathic properties?	0.090	0.00864
	Tolerates herbivory pressure?	0.472	0.0456
	Normal growth rate?	0.192	0.018432
	Stress tolerances?	0.185	0.01776
Reproduction		0.119	
	Reproductive system?	0.047	0.005593
	Propagule production?	0.460	0.05474
	Seed longevity?	0.256	0.030464
	Reproductive period?	0.101	0.012019
	Time to reproductive maturity?	0.136	0.016184
Dispersal		0.284	
	Number of mechanisms?	0.333	0.094572
	How far do propagules disperse?	0.667	0.189428

By comparing the major groups (i.e., establishment, growth/competitive ability, reproduction and dispersal), the working party determined the following order of importance of invasiveness indicators:


- 1) the plant's ability to establish in an ecosystem,
- 2) its ability to disperse,
- 3) its reproduction strategy, and
- 4) its growth/competitive ability.

These group weightings can also be expressed graphically (Figure 2). Within each group, the individual criterion were compared and weighted against each other (

Table 3). For instance, within the dispersal group, the working party decided that the question 'how far do propagules disperse?' was twice as important as the 'number of mechanisms' for dispersal (

Table 3). The results of the intra-group criteria weightings are also shown graphically (Figure 3).

Figure 2. Group weightings of invasiveness.



Within the invasiveness hierarchy, the weightings of individual criteria are multiplied by the group's weighting (eg. 'distance of dispersal' x 'dispersal' $\Rightarrow 0.667 \times 0.284 = 0.189$), to produce a total weighting for each criterion (Figure 4). During the assessment of biological data, criteria are assigned intensity ratings (criteria ratings) of high (H), medium-high (MH), medium (M), medium-low (ML), and low (L), to score each species. (The respective numerical value for each intensity rating is H=1, MH=0.67, M=0.5, ML=0.33 and L=0.) The scored intensity ratings for each criterion and their weightings are then tallied and calculated to produce a final 'invasiveness score' for each species, ie.

$$\text{Invasiveness score} = \sum ((\text{Group weighting} \times \text{Criterion weighting}) \times \text{Intensity rating})$$

The closer the final invasiveness score is to 1, the more invasive the plant is. The invasiveness score for each species is only relative to scores of other plants run through the same process, but can be used to rank species as to their potential invasiveness or rate of spread. An example of this process is shown for gorse/furze and boxthorn (Table 4). A summary of biological data was collated to determine the 'invasiveness score' of gorse/furze and boxthorn (Table 4). The information to rate each criterion was obtained from databases, journal articles, flora's of the world (books or articles describing the species of a particular country or region), online information, and other sources. There is much available information on some species (eg. declared noxious species), and very scant information for others (eg. grasses and new and emerging weed species). Where there is an information gap for a particular criterion, a 'medium' (M) ranking is given to indicate 'unknown'. Although the invasiveness assessments are undertaken using the best available information, they are only as accurate as the information that is used. Therefore, as we become more informed about a species, reassessment may be necessary.

The invasiveness criteria and intensity ratings are in Appendix 1.




Figure 3. Criteria weightings of invasiveness.




Figure 4. Total criteria weightings of invasiveness.

Table 4. Comparison of invasiveness assessments for gorse/furze *Ulex europaeus* and boxthorn *Lycium ferocissimum*.

QUESTION	<i>Ulex europaeus</i>		<i>Lycium ferocissimum</i>	
	COMMENTS	RATING	COMMENTS	RATING
Establishment				
Germination requirements?	“Most germination occurs in autumn and spring to mid summer”.	MH	Seeds germinate at any time of the year.	H
Establishment requirements?	Occurs in ecosystems receiving shading e.g. dry and damp sclerophyll forest.	MH	Occurs in open areas.	ML
How much disturbance is required?	Invades undisturbed ecosystems – heathland and heathy woodland.	H	Establishes in pastures. Listed as a ‘problematic weed of agricultural situations’	MH
Growth/Competitive				
Life form?	Fabaceae family: a legume.	MH	Large shrub, other category	L
Allelopathic properties?	No allelopathic properties described.	L	No allelopathic properties described.	L
Tolerates herb pressure?	“Young seedlings, which are not armed with stiff spines, are readily grazed by sheep and rabbits”.	MH	Seldom grazed by stock because of the sharp spines.	MH
Normal growth rate?	“Rapidly growing”. “Control is made difficult by the vigour and comprehensiveness of the plant”.	MH	Early root growth is rapid ensuring young plants are competitive	H
Stress tolerance to frost, drought, w/logg, sal. etc?	“Tolerates many soil types, frost to -20°C (young plants are sensitive), salt laden wind and drought”. Burns readily but not killed – re sprouts and seeds germinate”.	MH	Tolerant of drought and frost. Some water logging (occurs in creek beds and along streams and rivers).	MH
Reproduction				
Reproductive system	“Reproducing by seed”.	ML	Reproducing by seed.	ML
Number of propagules produced?	“Seed production is prolific with an annual input of up to 6 million seeds per hectare”.	H	Fruit production has ranged between 0535. Seeds numerous. ~ 500 fruit x 10 seeds per fruit=5,000 seeds.	H
Propagule longevity?	“Have a high viability and even seeds 25 years old is 85% viable”.	H	No information available.	M
Reproductive period?	“Plants are long lived, producing new	H	Large shrub. Produces dense thickets.	H

	growth each spring". "Living to a maximum life of 29 years".			
Time to reproductive maturity?	"Plants <u>may</u> flower first when about 18 months old".	MH	Plants do not flower until at least 2 years old.	MH
Dispersal				
Number of mechanisms?	Refer to 'dispersal' in P & C (1992 p. 482) → wind, birds, ant's etc.	H	The fruit is commonly eaten by birds and foxes and the seeds are viable when excreted.	H
How far do they disperse?	"Birds are important in spreading seeds and patches of weeds are often found under trees or posts where birds have perched".	H	Above animals could disperse seeds > 1 km as quite mobile.	H
Invasiveness Index (max. = 1, min. = 0)		0.83		0.67

1.5 Present and Potential Distribution of Pest Plants

Present and potential distributions are another major component required in the decision support system and AHP to predict the status of a weed. The greater the potential distribution of a weed the greater the potential impact and management costs. To ensure the most cost-effective use of weed management resources, invasive species that have the greatest potential range should be targeted. Prioritisation is also important as it is unrealistic to expect that all weeds can be controlled with limited available resources. Knowledge of potential distribution is also necessary for devising management programs. Land managers can be alerted to the risk of weed invasion and measures can be enforced to prevent the introduction of weed propagules into new areas. Low priority can be given to areas where the weed might fail to persist, or be of little economic, environmental or social importance.

In determining the potential distribution of plants, consideration must be given to the environmental conditions that a given genotype is exposed to. The US National Research Council for example, pointed out that;

" simply identifying the traits of a species and ignoring the environmental context in which the species occurs limits the information about whether the species can persist, let alone become invasive " (NRC 2002).

Two of the major environmental factors influencing weed distribution are climate and land-use or vegetation type. Weed species are typically more invasive in regions that are climatically similar to their native environment. Climate limits distribution according to how temperature and moisture stresses affect the weed's life cycle. Different land-uses (eg. cropping, perennial pasture and forestry) or vegetation types (eg. Grasslands, woodland, mallee) have different disturbance regimes that may or may not favour different groups of weeds. Therefore, having determined the climatic preferences of a weed, it is necessary to overlay its potential location on a map of the weed's associated land-use/vegetation types in Victoria. The areas of the state that are potentially at risk from this weed can then be identified.

1.6 Present Distribution

Information on the weed's present distribution, both overseas and in Australia, is collected from databases, journal articles, floras of the world (books or articles describing the species of a particular country or region), online information, and any other sources. Serrated tussock *Nassella trichotoma* (Nees) Hack. Ex Archav. is used here as an example to highlight the variety of sources, and the process, used to determine a weed's present distribution.

Serrated tussock originates from Argentina and surrounding countries (Chile, Peru and Uruguay) in South America (Figure 5). Serrated tussock now occurs in South Africa, New Zealand, occasionally in Europe (eg. Italy) (Figure 6), and in Australia (Figure 7) but has not been reported elsewhere. In Australia it is thought the weed was introduced about 1900, but not recorded until 1935, growing at Yass in NSW. It was first recorded in Victoria in 1954, and Tasmania in 1956, and does not occur in any of the other states. In NSW it is most widespread on the southern and central tablelands but also occurs widely on the New England tablelands. In Victoria it is mainly found on the basalt plains west of Melbourne (Figure 8). Smaller infestations occur in the Western District, in Gippsland, and in a small patch in the north east of the State (Figure 8). A small area of serrated tussock occurs in Tasmania, near Hobart, and on King Island in Bass Strait.



Figure 5. Distribution of serrated tussock in countries of origin.



Figure 6. Distribution of serrated tussock worldwide- except Australia.




Figure 7. Distribution of serrated tussock in Australia.




Figure 8. Known naturalisations of serrated tussock in Victoria (From DPI and DSE's IPMS).

1.7 Potential Distribution

Information on Australian and overseas distributions were imported into a climate-matching program, CLIMATE (Pheloung 1996), to predict potential distribution in Australia. Using the localities where a species occurs overseas and within Australia, the potential climatic range of any species can be overlaid upon Australia's climatic regions. The maps below illustrate the climatic regions most suitable for serrated tussock in Australia (Figure 9) and Victoria (Figure 10).




Figure 9. Potential distribution of serrated tussock in Australia, according to climatic parameters (Areas in red indicate a 80%+ match with the preferred climate of the plant species, yellow 70%, orange 60% and light green 50%).




Figure 10. Potential distribution of serrated tussock in Victoria, according to climatic parameters (Areas in red indicate a 80%+ match with the preferred climate of the plant species, yellow 70%, orange 60% and green 50%).

The 16 climatic parameters that are used to determine potential distribution can be grouped according to temperature or rainfall (Figure 11). Aquatic weeds are modelled for potential climatic range differently than terrestrial species. Rainfall is not a major criterion for determining the potential range of aquatic species, especially submergents, although it may play an important role in triggering certain biological properties (eg. freshwater floods appear to stimulate flowering in *Spartina*) (Strong pers. comm.). Thus rainfall parameters are excluded when predicting the climatic range of aquatic weeds. Water temperature is generally more moderate and

has fewer fluctuations than air temperature, and would provide a more accurate prediction for modelling aquatic species, however the necessary data is usually unknown. Therefore, modelling the climatic range of aquatic species has included eight air temperature parameters that provide at least some indication of potential range. The process for aquatic weeds is, consequently, more uncertain and likely to overestimate the species' actual potential range.




Figure 11. Dialogue box from CLIMATE (Pheloung 1996) showing the climatic parameters used in terrestrial weed modelling. The eight rainfall parameters are not included when modelling the potential climatic range of aquatic weeds.

Climatic overlays are then used to determine the potential range of the plant species by overlaying or intersecting them with susceptible land-uses, and broad vegetation types (BVTs) or wetlands, using ArcView GIS 3.2a software (ESRI Australia Pty Ltd, Melbourne). This refines potential distribution maps by recognising that plants are limited by factors other than climate alone, such as disturbance regimes associated with land-uses.

In the serrated tussock example, the weed is known to prefer sub-humid, subtropical and warm-temperate regions, and to occur as a weed in native and introduced pastures and lightly timbered regions. It is not restricted to any soil type or rainfall pattern and is relatively resistant to drought. In South America it is a minor weed invading cleared woodland, ploughed fields and neglected areas (Parsons and Cuthbertson, 1992). Serrated tussock also invades dry coastal vegetation, lowland grassland & grassy woodlands, dry sclerophyll forests & woodlands, and rock outcrop vegetation (Carr *et. al.* 1992). From this information one can then use a variety of GIS layers to determine susceptible land-use and broad vegetation types. The susceptible land-use overlays in this case were 'irrigated pasture' and 'dryland pasture', and the susceptible native vegetation corresponded to the following broad vegetation types;

Coastal scrubs and grasslands; dry foothill forests; grasslands; plains grassy woodlands; valley grassy forests; coastal grassy woodlands; montane grassy woodlands; riverine grassy woodlands; rainshadow woodlands; and box-ironbark forests.

The resulting map (Figure 12) illustrates the potential range of this weed in Victoria.




Figure 12. Potential distribution of serrated tussock in Victoria according to climatic parameters, susceptible broad vegetation types (BVT's), and susceptible land-uses. (Areas in red indicate a very high probability that serrated tussock could establish in agricultural or natural ecosystems within this region, yellow a high, orange a medium, and green a low probability of establishment. In the non-coloured areas the plant is unlikely to establish as the climate, or land-use/vegetation is not presently suitable.)

1.8 Ratio of Present to Potential Distribution

The ratio of present to potential distribution provides an indication as to the stage that spread of a weed has reached. Another way of expressing this is the relative position of the species on its invasion graph (Figure 13). Weeds that have reached, or nearly reached, the full limit of their distribution are not a major concern in terms of potential spread and impacts. However weeds currently occupying a small area of their potential range, which are in the 'lag' or 'sleeper' phase, should become a management priority. An important outcome of comparing present to potential distribution is the ability to target early intervention actions against weed invasions more effectively. Early intervention not only achieves better results from government and land manager investment (Figure 14), but also reduces costs of control and impact on social, environmental and economic values.

Intensity ratings for evaluating the ratio of present to potential distribution are shown in Table 5. Intensity ratings are subsequently "ground-truthed" to ensure the rating and corresponding descriptors reflect what officers of relevant agencies are seeing in the field.





Figure 13. Invasion graph indicating stages of expansion of a new species into a habitat. (Adapted from Groves (1992) and Hobbs (1991)).

Table 5. Intensity ratings for evaluating the present compared to potential distribution of a weed.

Rating	Weight	Pres:Pot Ratio	Statewide Descriptive	Regional or CMA Descriptive Rating
Very High	1.00		Infestation(s) that are able to be eradicated with no chance of reinvasion from outside of area of control (interstate.).	Infestation(s) that are able to be eradicated with no chance of reinvasion from outside of area of control (interstate/ other region etc.)
High	0.85	> 100,000	Infestation(s) that are able to be eradicated with some chance of reinvasion, less than 1,000ha in Vic.	Infestation(s) that are able to be eradicated with some chance of reinvasion
Medium High	0.71	> 1:10,000	Several or widely scattered small infestations or one large infestation	Several small infestations beyond eradication
Medium	0.57	1:100 - 1:10,000	Several large infestations or lots of multiple widely scattered infestations or a few combinations of both	A large partially dispersed infestation or few widely scattered small infestations
Medium Low	0.42	1:10 - 1:100	Multiple large infestations and multiple small infestations.	Numerous large dispersed infestations or lots of scattered small infestations.
Low	0.28	1:2 - 1:10	The majority of region infested with some large areas still "clean" (more "clean" areas than infested)	The majority of region infested with some large areas still "clean" (more "clean" areas than infested)
Very Low	0.14	< 1:2	The majority of region infested with some smallish areas still "clean" (less "clean" areas than infested)	The majority of region infested with some smallish areas still "clean" (less "clean" areas than infested)
Extremely Low	0	1:1	Reached full potential – but may increase in density within infested area or no suitable climate match across Victoria.	Reached full potential – but may increase in density within infested area or no suitable climate match within the CMA

Figure 14. Total cost of plant invasions showing costs of early expenditure (Area A) and the resulting benefit (Area B) (Adapted from Hobbs and Humphries (1995)).



1.9 Limitations of Present and Potential Distribution Maps

Potential distribution maps are estimates and are only as reliable as the data they are based on. As more records are collected on where plants occur the predictions will become more accurate. It is expected, consequently, that there are potential distribution maps that do not yet fully represent existing or potential distribution.

For some species there may be insufficient data to undertake potential distribution mapping. For other species, information on present distribution may be under-represented in the databases used, with the exception of priority weeds such as serrated tussock. Conversely, the modelled potential distribution of weeds is likely to be overestimated. This occurs as the broad scale (*i.e.* 1:250,000) of the statewide databases used merges minor differences into the larger BVT's or land-uses for each grid. Microhabitats within a vegetation or land-use type may be unsuitable for the particular weed species, and microhabitats outside the identified susceptible land-use or vegetation type may be suitable but not recognised (eg. roadsides, small riparian or vegetation corridors). More detailed map layers, such as the soon to be introduced Ecological Vegetation Divisions (EVDs), and updated land use data will produce predictions of finer detail.

The many weeds recorded as occurring along roadsides presents another major limitation when predicting potential distribution. Victoria has over 170,000 kilometres of roads, however to include all these roads within the image would not be suitable, as it would be too cluttered and meaningless. Thus, some potential distribution images may not include the occurrence of weeds within a region, if they only occur along roadsides. For example, horehound *Marrubium vulgare* L. can occur along roadsides within cropping regions, but is unable to withstand cultivation. Similarly, some riparian weeds may occur along small rivers, streams and water channels, but these watercourses are too small or scattered to be detected at a 1:250,000 scale. As they are not included in the riparian or riverine vegetation classes of the BVT GIS layer, they do not appear on the predicted potential distribution maps.

These limitations highlight the need for ongoing action to improve our knowledge of weed distribution. Where information on a weed's present distribution is known but not recorded, records need to be updated to ensure management and monitoring are effectively undertaken

1.10 Impacts

The next stage of the WRA, before calculating a weed risk score (Section 1.3), is to determine the social, environmental, and economic impacts of pest plants. Some impacts of weeds can be measured in economic terms. These include:

- lost agricultural and horticultural production,
- cost of control of weeds, including fuel reduction activities,
- loss of recreational and tourism income as a result of degraded natural attractions,
- loss of native flora and fauna with potential commercial uses, and
- loss of water quality because of altered hydrological cycles and degraded natural landscapes.

However impacts on many social and environmental values, such as the impact of weeds on cultural sites or biodiversity, can be difficult to measure in dollar terms. Consequently it is yet to be demonstrated that we can credibly evaluate the full effects of weeds in such terms. Alternative assessment procedures have used general questions such as “does the weed have major, positive or negative effects on environmental health” (Virtue, pers comm.). Other procedures have been relatively detailed but lacked the transparency of weightings of questions.

Developing a process to measure impacts of weeds

Specific criteria to assess impacts have now been developed and weighted using a process similar to that for invasiveness (see Section 1.4). The method used AHP to generate weightings for individual criteria by pair-wise comparisons of criteria within subgroups, and of subgroups, as per the methodology designed by Saaty (1995).

A series of workshops, and surveys of workshop participants, identified the social, environmental and economic values that are, or could be, threatened by weeds. A list of criteria was developed at these workshops, and twenty-six specific criteria were selected as being appropriate to determine the impacts that weeds have on these values in Victoria (Table 6).

The Australian Standard for National Post-Border Weed Risk Management Protocol (AS/NZS HB 294:2006 Standards Australia/Standards New Zealand, 2006) identifies six factors fundamental to determining the impact of weeds. The factors question the degree to which the weed could:

- reduce the establishment of desired plants?
- reduce the biomass/yield of desired plants?
- reduce the quality of products or services?
- restrict physical movement?
- affect human and/or animal health?
- ecological processes?

The impact criteria established for this process reflect those factors and, in fact, expand upon them.

The workshops grouped these criteria according to a basic hierarchy of social, environmental (abiotic and biotic), and economic (principally agricultural) issues. Finally the criteria within groups were allocated weights according to pair-wise comparisons. Subgroups within the same group were also compared and weighted (ie. Tourism versus Cultural, or Vegetation versus Fauna), as were the major groups; social values (social impacts), natural resources (environmental impacts-abiotic), flora and fauna (environmental impacts-biotic), and agriculture (economic impacts) (Table 6).

The process included a revision of preliminary weightings to identify inconsistencies between perceived values and the weightings that had been assigned. For example, most social criteria were assumed less important than environmental criteria, however criteria such as threatened flora, or high value EVCs were found to be under-weighted, while some of the social criteria were over-weighted. The revision and adjustment of weightings in this fashion is in accordance with Saaty (1995). Final weighted criteria are presented in Figure 15 and, according to importance, in Figure 16.

Once weightings are finalised, individual weeds are subsequently assessed for each criterion according to 'intensity ratings' (Table 7). Each criterion is rated on a common scale (ie. 1 to 0) as H, MH, M, ML, or L (with respective intensity score of 1, 0.67, 0.5, 0.33 and 0). This intensity rating score is then multiplied by the overall weighting for that particular criterion and summed with all the other criteria scores to produce an overall impact score from 1 (scored H for all criteria) to 0 (scored L for all criteria).

The calculation of the impact score can be expressed:

$$\text{Impact Score} = \Sigma ((\text{Criteria Group Weighting} \times \text{Criteria Weighting}) \times \text{Criteria Intensity Rating})$$

1.11 Calculating a weed risk score

The final stage of the WRA is to apply the results of invasiveness (Section 1.4), distribution (Section 1.5), and impact (Section 1.6) assessments to determine the relative importance of weeds by calculating a weed risk Score. The formula for calculating the weed risk score (introduced in Section 1.3) is:

$$\text{Weed risk score} = \alpha (\text{Invasiveness score}) + \beta (\text{Present:Potential Distribution}) + \delta (\text{Impact})$$

$$\text{where } \alpha = 0.12, \beta = 0.32 \text{ and } \delta = 0.56$$

The application of this formula can be illustrated using the example of serrated tussock in two different catchment management authority (CMA) regions:

Wimmera CMA

$$\begin{aligned} &\text{Weed Risk Score} \\ &= \alpha (0.7615) + \beta (0.85) + \delta (0.6259) \\ &= 0.12 (0.7615) + 0.32 (0.85) + 0.56 \\ &\quad (0.6259) \\ &= \mathbf{0.7139} \end{aligned}$$

Port Phillip and Westernport CMA

$$\begin{aligned} &\text{Weed Risk Score} \\ &= \alpha (0.7615) + \beta (0.28) + \delta (0.6259) \\ &= 0.12 (0.7615) + 0.32 (0.28) + 0.56 \\ &\quad (0.6259) \\ &= \mathbf{0.5315} \end{aligned}$$

In this example serrated tussock was subsequently ranked as the third highest priority weed in the Wimmera CMA (with a relatively high weed risk score of 0.7139). In the Wimmera CMA serrated tussock currently occupies a small area of its potential range, therefore intervention in this case would optimise government and land manager investment, and reduce costs of control, and impact on social, environmental and economic values. In the Port Phillip and Westernport CMA however, serrated tussock (with a weed risk score of 0.5315) already occupies a large proportion of its potential distribution and was consequently ranked 32 of 101 weeds assessed.

Table 6. Group and criteria ratings for determining impact.

GROUP	SUB GROUP	CRITERIA	GROUP WEIGHTING	SUB GROUP WEIGHTINGS	CRITERION WEIGHTING	FINAL CRITERION WEIGHTS
Social Values			0.1			
	Tourism			0.875		
		Restrict human access?			0.297	0.0259875
		Reduce the 'tourism / aesthetics/ recreational use of the land'?			0.539	0.0471625
		Plant injurious, toxic, or spines affect people?			0.164	0.01435
	Cultural	Damage to indigenous or European cultural sites?		0.125		0.0125
Natural Resources - soil, water and processes			0.25			
	Water			0.5		
		Impact on water flow within watercourses or water bodies?			0.333	0.041625
		Impact on water quality (ie. dissolved O ₂ , water temperature)?			0.667	0.083375
	Soil	Increase soil erosion?		0.3		0.075
	Processes			0.2		
		Reduce the biomass of the community? (nb. biomass acting as a carbon sink).			0.1	0.005
		Change the frequency or intensity of fires?			0.9	0.045
Flora and Fauna			0.425			
	Vegetation			0.65		
		EVCs: Impact on the vegetation composition on the following:			0.53	
		a. High value EVCs			0.56	0.081991
		b. Medium value EVCs			0.34	0.04978025
		c. Low value EVCs			0.1	0.01464125
		Structure of a vegetation community?			0.25	0.0690625
		Threatened flora spp.?			0.22	0.060775
	Fauna			0.35		
		Threatened fauna spp.?			0.368	0.05474
		Non-threatened fauna spp.?			0.177	0.02632875
		Benefits or facilitates the establishment of indigenous fauna?			0.155	0.02305625
		Toxic, its burrs or spines affect indigenous fauna?			0.112	0.01666
		Pest Animals			0.188	
		Provide a food source to assist in success of pest animals?			0.4	0.011186
		Provide important habitat or harbour for serious pests?			0.6	0.016779
Agriculture			0.225			
		Quantity or yield of agricultural produce?			0.084	0.0189
		Agricultural quality (eg. contamination)?			0.144	0.0324
		Affect land value?			0.243	0.054675
		Change in priority of land use?			0.448	0.1008
		Increases the cost of harvest?			0.053	0.011925
		Act as an alternative host or vector for diseases of agriculture?			0.028	0.0063




Figure 15. Total criteria weightings of impact.




Figure 16. Total criteria weightings of impact according to importance

Table 7. Example of an impact assessment for serrated tussock according to criteria and intensity ratings. H=1, MH=0.75, M=0.5, ML=0.25, L=0

Question	Comments	Rating
1. To what extent does the weed restrict human access?	“A perennial tussock-forming grass to 50cm high and 25 cm diameter at the base.” Leaf spread is about 50 cm. Intensity rating: Would not hinder human access. (P & C 2001, Groves <i>et al</i> 1995).	L
2. To what level does this weed reduce the ‘tourism / aesthetics/ recreational use of the land?’	Tussocks are noticeable, but would not restrict recreational activities. Intensity rating: Weeds not obvious to the average visitor.	L
3. To what level is the plant injurious, toxic, or spines affect people?	Not toxic to humans.	L
4. How much damage is done to indigenous or European cultural sites?	Dense patches of tussocks likely to create a negative visual impact on cultural sites.	ML
5. To what extent does this weed impact on water flow within watercourses or waterbodies?	Terrestrial sp. (P & C 2001)	L
6. To what extent does the weed impact on water quality (ie. dissolved O₂, water temperature)?	Terrestrial sp. (P & C 2001)	L
7. To what extent does the weed increase soil erosion?	Tussocks persist for many years. Roots are diffuse and fibrous, mostly in top 20 cm of soil, and even seedlings are difficult to pull from the soil. Not likely to contribute to soil erosion. Intensity rating: Low probability of large scale soil movement. (P & C 2001)	L
8. To what extent does this weed reduce the biomass of the community? (n.b. biomass acting as a carbon sink).	“It dominates pastures and invades natural areas forming dense swards.” Intensity rating: Replaces biomass. (Blood 2001)	ML
9. To what extent does the weed change the frequency or intensity of fires?	“ <i>N. trichotoma</i> burns readily in winter. A dense mature infestation of <i>N. trichotoma</i> generally has 5–20 tussocks per square metre.” “Dense stands produce a serious fire hazard.” CFA trials (at Melton 1998) have indicated serrated tussock burns with an intensity up to seven times greater than grasslands. Seed heads create additional hazards where they build up against housing, sheds, roadsides, fence lines etc. This is a particularly serious hazard in the rural/urban interface areas of outer west Melbourne. Intensity rating: Dense infestations would moderately change the frequency and intensity of fire risk. (Groves <i>et al</i> 1995, Eurobodalla Shire, David Boyle, pers. comm)	H

Question	Comments	Rating	
10. To what extent does this weed impact on the vegetation composition on the following:	(a) High value EVCs	EVC=Plains grassland (E); CMA=Corangamite; Bioreg=Victorian Volcanic Plain; VH CLIMATE potential. Prolific seeder. Each plant can cover a large area leading to almost complete cover in dense infestations and eliminating most other species. Serious impact on grasses/forbs. 3-fold effect: competitiveness, water usage high, and allelopathy. Can form a monoculture if the area of infestation is 20% or greater-cannot save grassland. Intensity rating: Potential for monoculture within a specific layer. (P & C 2001, Colin Hocking – Pers comm)	H
	(b) Medium value EVCs	EVC=Grassy dry forest (E); CMA=Port Phillip; Bioreg=Central Victorian Uplands; VH CLIMATE potential. Also grows in lightly timbered areas. Impact as in 10(a) above. (P & C 2001)	H
	(c) Low value EVCs	EVC=Heathy dry forest (E); CMA=Glenelg Hopkins; Bioreg=Goldfields; VH CLIMATE potential. Impact as in 10(b) above. (P & C 2001)	H
11. To what extent does this weed effect the structure of a vegetation community? <i>(How many levels within community would be affected; Total of 6 levels – trees > 20m, trees 10–20m, shrubs 2–10m, shrubs < 2m, tussock grasses, ground covers & herbs (after Specht, 1970)).</i>	“In dense stands foliage of <i>N. trichotoma</i> completely covers the soil surface, thereby suppressing competitors.” “Mature plants develop a drooping, smothering form eventually excluding other ground-flora.” <i>N. trichotoma</i> is most commonly found in grassland/pasture situations, where it would compete strongly with and possibly replace other grasses, and forbs. “...establishing dense tree and shrub cover will, after a number of years, significantly impede Serrated Tussock invasion.” Intensity rating: Affects the lower two levels only. (P & C 2001, Muyt 2001)	ML	
12. What effect does the weed have on threatened flora spp.?	From serrated tussock establishment it takes only seven years to dominate a pasture or native grassland. Effects on <i>Danthonia</i> spp. Threatens ANZECC rated rare or threatened native plant species. (Groves <i>et al</i> 2003)	H	
13. What effect does the weed have on threatened fauna spp.?	Hotter burns impact on striped legless lizard, and other ground dwelling species. Loss of flora biodiversity as serrated tussock displaces desired species impacts on adequate food supply. Golden sun moth, (<i>Synemon plana</i>) are dependant on <i>Austrodanthonia</i> spp., which are displaced by serrated tussock. Intensity rating: Habitat changed leading to possible extinction of a VROT or Bioregional Priority spp. (A review of the conservation status of selected Australian non-marine invertebrates. G Clarke F Spier 2004)	H	
14. What effect does the weed have on non-threatened fauna spp.?	“Plants are unpalatable and infestations commonly expand as other species are selectively grazed out. The smothering form eventually excludes other ground-flora.” Reduction in habitat for native fauna. Or habitat changed dramatically? Possible local extinction. Wombats foraging in Monaro plains – foraging limited by ST infestations – forced to raid rubbish bins. Intensity rating: Habitat changed dramatically, leading to possible extinction. (Muyt 2001, David Boyle pers comm)	H	
Q15. To what extent does this weed provide benefits or facilitates the establishment of indigenous fauna?	No documented benefits for fauna. Intensity rating: Provides very little support for desirable species.	H	

Question	Comments	Rating
16. To what extent is the plant toxic, its burrs or spines affect indigenous fauna?	“Sheep will not graze it unless forced to and they lose weight and die due to a rumen full of undigested leaves.” May have similar impact on fauna, although they are unlikely to graze. Intensity rating: May cause fauna to lose condition. (Blood 2001)	ML
17. To what extent does this weed provide a food source to assist in success of pest animals?	Not known as a food source to pests.	L
18. To what extent does this weed provide important habitat or harbour for serious pests (eg. foxes, rabbits)?	Across the basalt plains rabbits are the greatest vertebrate environmental pest and serrated tussock provides harbour; permanent harbour in Rowsley Valley area. Intensity rating: Capacity to provide harbour for rabbits throughout the year. (David Boyle pers comm)	H
19. To what extent does this weed impact on the quantity or yield of agricultural produce?	“Even moderate infestations reduce carrying capacity by about 40% and up to 100%.” Serious impact on quantity of produce. Intensity rating: Serious impact on quantity e.g., >20% reduction. (P & C 2001, David Boyle pers comm)	H
20. To what extent does the weed impact on agricultural quality (eg contamination – lower price)?	Animals forced to eat the plant lose condition. “Even a moderate loss of condition results in lowered wool quality because of loss of crimp and breaks in the fleece. Seeds also contribute to vegetable fault in wool.” Major impact on quality. Canola crop contaminated by seed contaminant by wind, civil court case - \$60,000 damage. Intensity rating: Serious impacts on quality. Produce rejected for sale or export. (P & C 2001, David Boyle pers comm)	H
21. To what extent does this weed affect land value?	As a serious weed of pasture with significant impact on carrying capacity and reduction in agricultural return, its presence would seriously affect land value. Seven (7) Local Govt councils have introduced rebate schemes to arrest the decline in land value. Intensity rating: Major significance > 10% reduction in land value. (P & C 2001, David Boyle pers comm)	H
22. To what extent does this weed cause a change in priority of land use?	In New Zealand, “considerable effort to control [<i>N. trichotoma</i>], involving government purchase of heavily infested farms, clearing the weed at government expense, and reselling the land for farming, has been expended over many years.” Without government intervention, the land would have had no use for pastoral activities. Intensity rating: Significant loss of land for agricultural use. (P & C 2001)	H
23. To what extent the presence of the weed increases the cost of harvest.	Not a weed of cropping in Australia. In uncultivated areas serrated tussock dominates but can be controlled with cropping regimes. In very heavy infestation areas continuous cropping is the only control option and creates another set of problems with soil health. However the impact of seeding tussock blown onto pre-harvested crops has resulted in downgraded quality of produce and has seen one civil case against neighbouring polluters. More seed testing carried out. Time taking in harvest and post harvest testing. Also need to cultivate each year to prevent re-establishment. Intensity rating: Major increase in time or labour, or machinery in harvesting. (David Boyle pers comm)	H
24. To what extent does this weed act as an alternative host or vector for diseases of agriculture?	None evident. Unknown.	M

Methodology References

- AS/NZS HB 294:2006 Standards Australia/Standards New Zealand. (2006) National post-border weed risk management protocol. Standards Australia, Standards New Zealand, CRC for Australian Weed Management, Sydney, NSW, Australia.
- Carr, G.W., Yugovic, J.V. and Robinson, K.E. (1992) *Environmental Weed Invasions in Victoria: Conservation and Management Implications*. Department of Conservation and Natural Resources and Ecological Horticulture, Victoria.
- Groves, R.H. (1992) *Weed ecology and spread*. In: Proceedings of the First International Weed Control Congress (Volume 1). Weed Science Society of Victoria, Melbourne.
- Hobbs, R.J. (1991) *Disturbance a precursor to weed invasion in native vegetation*. Plant Protection Quarterly 6(3): 99-104.
- Hobbs, R.J. and Humphries, S.E. (1995) *An Integrated Approach to the Ecology and Management of Plant Invasions*. Conservation Biology 9(4): 761-770.
- NRC (National Research Council), Committee on the scientific basis for predicting the invasive potential of non-indigenous plants and plant pests in United States. (2002) *Predicting invasions of non-indigenous plants and plant pests*. National Academy Press, Washington
- Pheloung, P.C. (1996) *CLIMATE: a system to predict the distribution of an organism based on climate preferences*. Agriculture Western Australia, Perth.
- Robertson M.P., Villet M.H., Fairbanks D.H.K., Henderson L., Higgins S.I., Hoffmann J.H., Le Maitre D.C., Palmer A.R., Riggs I., Shackleton C.M. and H.G. Zimmermann. (2003), *A proposed prioritization system for the management of invasive alien plants in South Africa*, South African Journal of Science 99: 1-7
- Ross, J. H. and Walsh, N. G. (2003) *A census of the vascular plants of Victoria*. 7th edition. National Herbarium of Victoria, Royal Botanic Gardens, Melbourne
- Saaty, T. (1995) *Decision Making for Leaders: The Analytical Hierarchy Process for Decisions in a Complex World*. RWS Publications, Pittsburgh.
- Sax, D.F. and Brown, J.H. (2000) *The paradox of invasion*. Global Ecology & Biogeography 9, 363–371
- Sposito, V., Thatcher, A., Itami, B., Weiss, J., and Cotter, M. (2002) *Priority setting framework for natural resources management - Application of the analytical hierarchy process and natural resources accounting*. Department of Natural Resources and Environment, Melbourne
- Williamson, M. and Fitter, A., (1996) *The varying success of invaders*. Ecology 77: 1661-1666

Appendix 1. Invasiveness criteria and intensity ratings

Criteria	Intensity Rating			
	Low	Medium Low	Medium High	High
Establishment				
1. Germination /propagule requirements?	Requires specific environmental factors that are not part of an annual cycle to germinate (eg. specific temperatures, or human caused disturbance, such as ploughing).	Requires unseasonal or uncommon natural events for germination (eg. flooding, fire).	Requires natural seasonal disturbances such as seasonal rainfall, spring/summer temperatures for germination.	Opportunistic germinator, can germinate or strike/ set root at any time whenever water is available.
2. Seedling/ propagule establishment requirements (<i>i.e.</i> light, water, nutrients)?	Requires additional and very specific factors such as nutrients and water that are deliberately added or highly eutrophic conditions.	Requires more specific requirements to establish (eg. open space or bare ground with access to light and direct rainfall).	Can establish under moderate canopy/litter cover	Can establish without additional factors.
3. How much disturbance is required for seedling establishment to occur?	Major disturbance required with little OR no competition from other plant species.	Establishes in highly disturbed natural ecosystems (eg. roadsides, wildlife corridors, or areas which have a greater impact by humans such as tourist areas or campsites) OR in overgrazed pastures/poorly growing or patchy crops.	Establishes in relatively intact OR only minor disturbed natural ecosystems (eg. wetlands, riparian, riverine, grasslands, open woodlands); in vigorously growing crops OR in well-established pastures.	Establishes in healthy AND undisturbed natural ecosystems (eg. mallee, alpine, heathland).

Criteria	Intensity Rating			
	Low	Medium Low	Medium High	High
Lifeform and competitive ability				
4. Life form?	Other.	Geophyte, climber or creeper.	Grass, leguminous plant.	Aquatic (submerged, emergent, floating for ALL of life, inc. germination), and semi aquatic (some plant parts always in water).
5. Allelopathic properties? *	None.	Minor properties.	Allelopathic properties seriously affecting SOME plants.	Major allelopathic properties inhibiting ALL other plants.
6. Ability to tolerate herbivory pressure and produce propagules?	Preferred food of herbivores. Eliminated by moderate herbivory or reproduction entirely prevented.	Consumed and recovers slowly. Reproduction strongly inhibited by herbivory but still capable of vegetative propagule production (by rhizomes or tubers); weed may still persist.	Consumed but non-preferred OR consumed but recovers quickly; capable of flowering /seed production under moderate herbivory pressure (where moderate = normal; not overstocking or heavy grazing).	Favoured by heavy grazing pressure as not eaten by animals/insects and not under a biological control program in Australia/New Zealand.
7. Normal growth rate?	Slow growth; will be exceeded by many other species.	Maximum growth rate less than, many species of the same life form.	Moderately rapid growth that will equal competitive species of the same life form.	Rapid growth rate that will exceed most other species of the same life form.
		Medium		
		Growth rate equal to the same life form, OR there is widely conflicting evidence.		

Criteria	Intensity Rating			
	Low	Medium Low	Medium High	High
8. Stress tolerance of established plants to frost, drought, water logging, salinity, fire?	Maybe tolerant of one stress, susceptible to at least two.	Tolerant to at least two AND susceptible to at least one.	Highly tolerant of at least two of drought, frost, fire, waterlogging, and salinity, AND MAY be tolerant of another. Susceptible to at least one.	Highly resistant to at least two of drought, frost, fire, waterlogging, and salinity not susceptible to more than one (cannot be drought or waterlogging).
Reproduction				
9. Reproductive system?	Sexual (either cross OR self-pollination).	Sexual (self AND cross-pollination).	Vegetative reproduction (may be via cultivation, but not propagation).	Both vegetative AND sexual reproduction (vegetative reproduction may be via cultivation, but not propagation).
10. Number of propagules produced per flowering event?	Less than 50.	50-1000.	1000-2000.	Above 2000.
11. Propagule longevity?	Greater than 25% of seeds survive 5 years, OR vegetatively reproduces.	Greater than 25% of seeds survive 5-10 years in the soil, OR lower viability but survive 10-20 years.	Greater than 25% of seeds survive 10-20 years in the soil, OR lower viability but survives over 20 years.	Greater than 25% of seeds can survive over 20 years in the soil.
12. Reproductive period?	Mature plant produces viable propagules for only 1 year.	Mature plant produces viable propagules for only 1–2 years.	Mature plant produces viable propagules for 3–10 years.	Mature plant produces viable propagules for 10 years or more, OR species forms self-sustaining monocultures.

Criteria	Intensity Rating			
	Low	Medium Low	Medium High	High
13. Time to reach reproductive maturity?	Greater than 5 years to reach sexual maturity, OR for vegetative propagules to become separate individuals.	2-5 years to reach sexual maturity, OR for vegetative propagules to become separate individuals.	Produces propagules between 1-2 years after germination, OR vegetative propagules become separate individuals between 1-2 years.	Reaches maturity and produces viable propagules, OR vegetative propagules become separate individuals, in under a year.
Dispersal				
14. Number of dispersal mechanisms?	Propagules mainly spread by gravity.	Deliberate human dispersal (propagation or planting).	Propagules spread by wind, water, attachment (humans, animals, or vehicles), OR accidental human dispersal (ploughing).	Very light, wind dispersed seeds, OR bird dispersed seeds, OR has edible fruit that is readily eaten by highly mobile animals.
15. Probability (or chance) that propagules will disperse to a distance greater than one kilometre?	Very unlikely to disperse greater than 200 metres, most less than 20 metres.	Very few to none will disperse to one kilometre, most 20-200 metres.	Few propagules will disperse greater than one kilometre but many will reach 200-1,000 metres.	Very likely that at least one propagule will disperse greater one kilometre.

* **Allelopathic** - definition: Release of a substance by a plant that inhibits the germination, or growth, of another plant. Release can be through glands, residue, root exudate, or volatilisation. From DiTomaso, J and Healy, E. (2003) *Aquatic and Riparian Weeds of the West*.

For question 5 (*allelopathic qualities*): if lots of info' about the species is available in literature but mostly pre-1984, **AND/OR** no mention in the scant literature after 1984; the default rating is L, with a confidence level of L. If lots of research literature post-1984 is available with no mention of allelopathy, the assessor MAY infer a rating of L, with a confidence rating of L.

Appendix 2. Impact criteria and intensity ratings

Criteria	Intensity Ratings			
	Lowest Threat L	ML	MH	Highest Threat H
SOCIAL (Tourism, Visual aesthetics, Experience, Cultural sites)				
1. To what extent does the weed restrict human access?	Minimal OR negligible impact (<i>ie.</i> can go anywhere).	Low nuisance value. Impedes individual access; unable to walk to waterways.	High nuisance value. People AND/OR vehicles access with difficulty.	Major impediment to access waterways OR machinery. Significant works required to provide reasonable access, tracks closed or impassable.
2. To what level does this weed reduce the 'tourism / aesthetics/ recreational use of the land'?	Weeds not obvious to the 'average' visitor.	Minor effects to aesthetics AND/OR recreational uses (<i>ie.</i> aware but not bothered or activity inhibited).	Some recreational uses affected.	Major impact on recreation. Weeds obvious to most visitors, with visitor response complaints AND a major reduction in visitors.
3. To what level is the plant injurious, toxic, or spines affect people?	No effect, no prickles, no injuries.	Mildly toxic, may cause some physiological issues (e.g. hayfever, minor rashes, minor damage from spines and burrs at certain times of year).	Spines, burrs or toxic properties at most times of the year, OR may be a major component in allergies, hayfever and/or asthma.	Large spines or burrs, extremely toxic, AND/OR cause serious allergies to humans throughout year.
4. How much damage is done to indigenous or european heritage sites, and infrastructure?	Little or negligible effect on aesthetics or structure of site.	Moderate visual effect.	Moderate structural effect.	Major structural damage to site, AND/OR obliteration of the heritage/cultural feature.
NATURAL RESOURCES – SOIL, WATER & PROCESSES				
5. To what extent does this weed impact on water flow within watercourses or waterbodies?	Little or negligible affect on water flow.	Minor impact on surface or subsurface flow either by roots or free floating aquatics.	Major impact on either surface OR subsurface flow (eg. major root obstructions, submergent aquatics).	Serious impacts BOTH to surface AND subsurface water flow (eg. attached emergent aquatics).

Criteria	Intensity Ratings			
	Lowest Threat L	ML	MH	Highest Threat H
6. To what extent does the weed impact on water quality (ie. dissolved O ₂ , water temperature)?	No noticeable effect on dissolved O ₂ OR light levels.	Noticeable but minor effects in either dissolved O ₂ OR light levels.	Noticeable but moderate effects in both dissolved O ₂ AND light; causing increased algal growth.	High effects in either dissolved O ₂ AND/OR light; causing eutrophication.
7. To what extent does the weed effect soil erosion?	Low probability of large scale soil movement; OR decreases the probability of soil erosion.	Moderate probability of large scale soil movement.	High probability of large scale soil movement with minor off-site implications.	High probability of large scale soil movement with major off-site implications.
8. To what extent does this weed reduce the biomass of the community? (nb. biomass acting as a carbon sink).	Biomass may increase.	Direct replacement of biomass by invader.	Biomass slightly decreased.	Biomass significantly decreased (eg. trees replaced by more open community).
9. To what extent does the weed change the frequency or intensity of fires?	Small or negligible effect on fire risk.	Minor change to either frequency OR intensity of fire risk.	Moderate change to both frequency and intensity of fire risk.	Greatly changes the frequency AND/OR intensity of fire risk.
FLORA & FAUNA/ VEGETATION & EVCS				
10. To what extent does this weed impact on the vegetation composition on the following:	Very little displacement of any indigenous spp. Sparse/ scattered infestations.	Minor displacement of some dominant or indicator spp. within any one strata/layer (eg. ground cover, forbs, shrubs & trees).	Major displacement of some dominant spp. within a strata/layer (or some dominant spp. within different layers).	Monoculture within a specific layer; displaces all spp. within a strata/layer.
a. High value EVC				
b. Medium value EVCs	(as above)	(as above)	(as above)	(as above)
c. Low value EVCs	(as above)	(as above)	(as above)	(as above)

Criteria	Intensity Ratings			
	Lowest Threat L	ML	MH	Highest Threat H
11. To what extent does this weed effect the structure of a vegetation community?	Minor or negligible effect on <20% of the floral strata/layers present; usually only affecting one of the strata.	Minor effect on 20-60% of the floral strata.	Minor effect on >60% of the layers or major effect on < 60% of the floral strata.	Major effects on all layers. Forms monoculture; no other strata/layers present.
12. What effect does the weed have on threatened flora spp.?	Minor/negligible effects on any Bioregional Priority or VROT spp.	Any population of a VROT spp is reduced.	Any population of Bioregional Priority 1A* spp is reduced, or any population of a VROT spp is replaced.	Any population of Bioregional Priority 1A* spp is replaced.
FLORA & FAUNA				
13. What effect does the weed have on threatened fauna spp.?	No threatened fauna affected due to fauna not co-existing within infested area or strata.	Minor effects on threatened spp.; minor hazard OR reduction in habitat/food/ shelter.	Reduction in habitat for threatened spp, leading to reduction in numbers of individuals, but NOT to local extinction.	Habitat changed dramatically, leading to the possible extinction (extirpation) of a VROT or Bioregional Priority spp.
14. What effect does the weed have on non-threatened fauna spp.?	No fauna affected due to fauna not co-existing within weed area or strata.	Minor effects on fauna spp.; minor hazard OR reduction in habitat/food/ shelter.	Reduction in habitat for fauna spp., leading to reduction in numbers of individuals, but NOT to local extinction.	Habitat changed dramatically, leading to the possible extinction (extirpation) of non-threatened fauna.
15. To what extent does this weed provide benefits, or facilitates the establishment of, indigenous fauna?	Provides vital food, shelter OR assists the recolonisation of desirable species.	Provides an important alternative food source and/or harbor to desirable species.	Provides some assistance in either food or shelter to desirable species.	Provides very little support to desirable species.
16. To what extent is the plant toxic, its burrs or spines affect indigenous fauna?	No effect.	Mildly toxic, may cause fauna to lose condition.	Spines, burrs or toxic properties to fauna at certain times of the year.	Large spines or burrs dangerous to fauna. Toxic, and/or causes allergies.

Criteria	Intensity Ratings			
	Lowest Threat L	ML	MH	Highest Threat H
FLORA AND FAUNA/ Pest Animal				
17. To what extent does this weed provide a food source to assist in success of pest animals?	Provides minimal food for pest animals.	Supplies food for one or more minor pest spp. (eg. blackbirds or environmental insect pests).	Supplies food serious pest (eg. rabbits and foxes), but at low levels (eg. foliage).	Supplies food for > 1 major pest spp at crucial times of the year (eg. heavy berry load or continual food throughout the year).
18. To what extent does this weed provide important habitat or harbor for serious pests (<i>pests for which DPI has a statewide program</i> eg. foxes, rabbits, fire ants,)?	No harbour for pest spp.	Doesn't provide harbor for serious pest spp, but may provide for minor pest spp.	Capacity to harbor rabbits or foxes at low densities or as overnight cover.	Capacity to provide harbor and permanent warrens for foxes and rabbits throughout the year.
AGRICULTURE – Quality, Quantity, Cost to Production, Effect on land use and value				
19. To what extent does this weed impact on the quantity or yield of agricultural produce?	Little or negligible affect on quantity of yield.	Minor impact on quantity of produce (eg < 5% reduction).	Major impact on quantity of produce (eg 5-20%).	Serious impacts on quantity (eg >20% reduction). Unviable to harvest crop/ stock.
20. To what extent does the weed impact on agricultural quality (eg. contamination – lower price)?	Little or negligible affect on quality of yield.	Minor impact on quality of produce (eg < 5% reduction).	Major impact on quality of produce (eg 5-20%).	Serious impacts on quality (eg >20% reduction). – Produce rejected for sale or export.
21. To what extent does this weed affect land value?	Little or none.	Decreases in land value <10%.		Major significance > 10%.
22. To what extent does this weed cause a change in priority of land use?	Little or no change	Some change, but no serious alteration of either agricultural return. Affects more the visual rather than intrinsic agricultural value.	Downgrading of the priority land use, to one with either less agricultural return.	Major detrimental change and significant loss for agricultural usage (eg complete change to different ag use eg farm forestry.)

Criteria	Intensity Ratings			
	Lowest Threat L	ML	MH	Highest Threat H
23. To what extent the presence of the weed increases the cost of production?	Little or none.	Minor increase in cost of harvesting – eg slightly more time or labour is required.		Major increase in time or labour, or machinery in harvesting.
24. To what extent does this weed act as an alternative host or vector for diseases of agriculture?	Little or no host.	Provides host to minor (or common) pests, or diseases.		Host to major and severe disease or pest of important agricultural produce.

** Information on bioregional priorities can be found in Bioregional Strategic Overviews. As at June 2005, not all Victorian Bioregions have been reviewed.*