

# **A STUDY OF THE LAND IN THE CATCHMENT OF LAKE EILDON**

ALLEN S. RUNDLE

**SOIL CONSERVATION AUTHORITY  
VICTORIA**

**1977**

\*Previously Soil Conservation Authority, now School of Earth Sciences, Macquarie University,  
Eastwood, N.S.W.

## SOIL CONSERVATION AUTHORITY



Chairman:

A. Mitchell, M. Agr. Sc., D.D.A

Deputy Chairman:

G. L. SWARTZ, M. Agr. Sc.

Member:

J. S. GILMORE

Secretary:

D.G. MORTON

## TABLE OF CONENTS

FOREWORD.....	v
SUMMARY .....	vi
PURPOSE AND AREA OF SURVEY .....	1
THE ENVIRONMENT OF THE CATCHMENT.....	2
GEOLOGY AND GEOMORPHOLOGY .....	2
Drainage .....	5
CLIMATE .....	6
Precipitation.....	6
Rainfall.....	7
Seasonal Rainfall.....	8
Effective Rainfall.....	8
Snowfall.....	8
Mist, Dew and Frost.....	8
Temperature.....	8
Humidity; Evaporation; Transpiration.....	12
Wind; Thunderstorms; Fire Danger Weather .....	12
SOILS.....	17
NATIVE VEGETATION.....	33
Classification .....	33
Patterns of Vegetation and the Factors Involved .....	34
Summary .....	46
THE LAND SYSTEMS .....	48
PRESENT USE AND CONDITION OF THE CATCHMENT .....	73
PRESENT LAND USE AND ITS HAZARDS .....	73
Kind of Production or Function.....	73
Major Management Practices and Hazards .....	76
CONDITION OF THE LAND .....	80
Erosion.....	80
Fire History.....	83
POTENTIAL USE OF THE CATCHMENT .....	85
ACKNOWLEDGEMENTS.....	89
REFERENCES .....	90

## INDEX TO TABLES

Table 1 – Stratigraphy and Structure of Lake Eildon Catchment .....	3
Table 2 – Predicted Mean Monthly Temperatures. (°C) .....	10
Table 3 – Soil Classification.....	15
Table 4 – Classification of the Plant Communities .....	35
Table 5 – Storage Characteristics of Eildon and Hume Catchments .....	73
Table 6 – Rating of Land Use Potential of Land Systems .....	85
Table 7. Estimated Proportion of the Precipitation which becomes runoff for each Land System (based on monthly precipitation, calculated evapotranspiration, soil depth and available water capacity) .....	86
Table 8 – Proportion of Inflow to Lake Eildon Contributed by each Land system (combines runoff proportion, precipitation and area of each land system).....	86

## INDEX TO FIGURES

Figure 1 – Inferred Precipitation .....	7
Figure 2 – Seasonal Rainfall.....	9
Figure 3 – Seasonal Temperature .....	11
Figure 4 – Calculated Seasonal Evapotranspiration at Various Altitudes .....	13
Figure 5 – Seasonal Water Budget at Mansfield, Jamieson and Woods Point .....	14
Figure 6 – Analytical Data for Solodic Soils, Gilgai Hummock (Profiles 315, 318) .....	19
Figure 7 – Analytical Data for Solodic Soils, Gilgai Hollow (Profiles 316, 317) .....	20
Figure 8 – Analytical Data for Chernozemic Soils (Profiles 319, 321).....	23
Figure 9 – Analytical Data for Yellow Podzolic Soil (Profile 329).....	25

Figure 10 – Analytical Data for Leptopodzols (Profiles 312, 313, 314).....	26
Figure 11 – Analytical Data for Amphipodzols (Profiles 320, 324).....	28
Figure 12 – Analytical Data for Acid Brown Earths (Profile 328).....	29
Figure 13 – Analytical Data for Alpine Humus Soils (Profiles 327, 330).....	30
Figure 14 – Relationships of dominant Vegetation in North-Eastern Victoria.....	40
Figure 15 – Relationships between Eucalypt Species, Rainfall and Altitude.....	41
Figure 16 – Mansfield land system.....	50
Figure 17 – Eildon land system.....	53
Figure 18 – Maintongoon land system.....	56
Figure 19 – Jamieson land system.....	59
Figure 20 – Howqua land system.....	62
Figure 21 – Bindaree land system.....	66
Figure 22 – Taponga land system.....	67
Figure 23 – Buller land system.....	72
Figure 24 – Precipitation and Water Production from Land System.....	87

## INDEX TO PLATES

Plate 1 – Lake Eildon, amongst its wooded hills.....	iv
Plate 2 – Lake Eildon has been formed by the construction of an earth and rock dam.....	1
Plate 3 – Model showing topography of the Eildon Catchment.....	2
Plate 4 – The Mansfield Plain.....	4
Plates 5 and 6 – Steep, rugged country characterises much of the Eildon catchment.....	4
Plate 7 – Gilgaied solodic soils of the Mansfield Plain.....	18
Plate 8 – A prairie soil overlies a buried chernozemic soil which lies on Quaternary gravels.....	21
of the Mansfield Plain.....	21
Plate 9 – Yellow and red podzolic soils in the Jamieson land system.....	22
Plate 10 – Acid brown earths are friable, porous, crumbly soils in the narrow-leaved peppermint ...	31
and alpine ash zones.....	31
Plate 11 – Young red gums in the Mansfield land system.....	34
Plate 12 – Forests of stringybark, box, peppermint and gum occur in the Maintongoon and adjacent	
land systems.....	34
Plates 13 and 14 – Peppermint gum forests characterise the extensive Howqua and Jamieson land	
systems.....	37
Plate 15 – Sub-alpine herbfields and grasslands are found in the cold air drainage basins of the	
Bindaree, Taponga and Buller land systems.....	38
Plate 16 – Coppice woodlands of snow gum are a feature of the Bindaree and Buller land systems.....	39
Plate 17 – Alpine herbfields and alpine heaths grow above the tree line in the Buller land system.....	39
Plate 18 – Aspect and seepage markedly control the distribution of the drier stringybark, box-	
peppermint forest and the wetter peppermint gum forest.....	44
Plate 19 – Mansfield land system, with hills of Eildon land system in distance.....	48
Plate 20 – Hilly part of Mansfield land system, with Eildon and Maintongoon land systems in	
distance.....	49
Plate 21 – Cleared hills of the Eildon land system surround Lake Eildon.....	51
Plate 22 – Timbered hills of the Maintongoon land system rise above the cleared slopes.....	54
of the Eildon land system.....	54
Plate 23 – The Jamieson land system is steeply sloping and covered with forests of.....	57
broad-leaved peppermint and gum.....	57
Plate 24 – Valleys in the Jamieson and Howqua land systems carry forests of candle gum.....	58
and narrow-leaved peppermint.....	58
Plate 25 – The Howqua land system ranges with long, steep slopes covered with forests of.....	60
narrow-leaved peppermint and gum.....	60
Plate 26 – Forests of alpine ash with mountain gum are found in the Bindaree and Taponga land	
systems. They are snow covered during winter.....	63
Plate 27 – Sub-alpine woodlands of snow gum are a feature of the Buller land system.....	68
Plate 28 – Grassy plains are found in the Buller land system where cold air drainage is ponded and	
prevents the establishment of snow gums.....	69
Plate 29 – The highest parts and exposed sites of the Buller land system extend into the treeless	
alpine zone, as in the foreground.....	70

Plate 30 – Snow melt in the high country is a very important component of the inflow into Lake Eildon. This picture also illustrates that dark surface absorb radiation and hasten the melting of snow .....	74
Plate 31 – Surge flows of muddy water should be avoided if possible.....	78
Plate 32 – Turbidity of water is aggravated by slumping of steep road batters. The natural flow of water is diverted and accelerated by roads .....	79
Plates 33 and 34 – Stream bank erosion in the Mansfield and Eildon land systems have been a source of silt to Lake Eildon .....	82
Plate 35 – Silt on the bed of Lake Eildon in 1962 .....	83
Plates 36 and 27 – Sub-alpine and alpine herbfields and grasslands are easily damaged, leading to bare spaces and decline in porosity. Recovery may be slow .....	84

### **LIST OF LAND SYSTEMS**

MANSFIELD LAND SYSTEM.....	48
EILDON LAND SYSTEM .....	51
MAINTONGOON LAND SYSTEM.....	54
JAMIESON LAND SYSTEM.....	57
HOWQUA LAND SYSTEM .....	60
BINDAREE AND TAPONGA LAND SYSTEMS .....	63
BULLER LAND SYSTEM.....	68

### **APPENDICES**

APPENDIX 1 - METHODS OF SOIL ANALYSIS .....	91
---	----