

Soil landform unit: MGn; Mountains on gneiss


Landform pattern	Mountains and some very high hills
Elevation (m)	300-920
Annual Rainfall (mm)	650-1500
Geomorphic unit	1.3.1 Eastern Uplands; Ridges, Valleys and Hills; High Dissected Ridges and Valleys
Geology	Ordovician gneiss and gneissic pegmatite



Locality diagram



Position in landscape



Landform component	1	2	3	4
Landform element	Crests and side slopes	Steep upper slopes	Moderate mid slopes	Moderate slopes in lower landscape position
Slope range (%)	5-40	20->32	20-32	10-20
Site drainage	Well to rapidly drained with high permeability	Moderately well drained with moderate permeability	Moderately well drained with moderate permeability	Moderately well drained with moderate to slow permeability
Rock outcrop	Mostly <2%, with rock size to 1 m	0-50%, rock to 0.6 m	10-20%, rock to 2 m	1%
Soil description	Yellowish brown to brown loam to light clay texture contrast soils	Yellowish brown to brown loam uniform soils	Reddish brown gradational and texture contrast soils	Brown loam to clay texture contrast soils
Present land use	Uncleared	Uncleared, grazing	Uncleared, grazing	Uncleared, grazing
Susceptibility to erosion	Sheet and rill: Moderate Gully: Moderate Landslip: High Wind: High	Sheet and rill: Moderate Gully: Moderate Landslip: High Wind: High	Sheet and rill: Moderate Gully: Moderate Landslip: Moderate Wind: Moderate	Sheet and rill: Moderate Gully: Moderate Landslip: Moderate Wind: Moderate
Native vegetation	<i>Eucalyptus goniocalyx</i> , <i>E. microcarpa</i> , <i>E. albens</i>	<i>E. goniocalyx</i> , <i>E. microcarpa</i> , <i>E. dives</i> , <i>E. macrorhyncha</i>	<i>E. goniocalyx</i> , <i>E. mannifera</i> , <i>E. dives</i> , <i>E. macrorhyncha</i>	<i>E. goniocalyx</i> , <i>E. mannifera</i> , <i>E. dives</i> , <i>E. macrorhyncha</i>